


AMENDMENTS TO THE CLAIMS:

The following listing of claims replaces all prior listings, and all prior versions, of claims in the application:

LISTING OF CLAIMS:

1. (Currently amended) A method of fluorination which comprises fluorinating a monosaccharide or monosaccharide bonded to a base of a nucleic acid using a fluorinating agent represented by general formula (I):




(I)

wherein Y represents nitrogen atom ~~or phosphorus atom~~, R^0 , R^1 and R^2 represent hydrogen atom or an alkyl ~~or aryl~~ group having 1 to 4 carbon atoms, and which may have substituents, ~~the atom and the groups represented by R^0 , R^1 and R^2 may be a same with or different from each other, and two or three of the groups represented by R^0 , R^1 and R^2 may be bonded to each other to form a ring, and R^0 represents a benzene ring or a pyridine ring, which has one or two substituents selected from the group consisting of alkyl group having 1 to 2 carbon atoms, halogen atom, methoxy group, amino group and nitro group, which may be the same as or different from each other.~~

2. (Original) A method of fluorination according to Claim 1, wherein, in general formula (I), Y represents nitrogen atom, R^0 represents 3-methyphenyl group or 2-methoxyphenyl group, and R^1 and R^2 represent ethyl group.


3. (Previously presented) A method of fluorination according to Claim 1, wherein the monosaccharide or monosaccharide bonded to a base of a nucleic acid is fluorinated by a thermal reaction.

4. (Currently amended) A method of fluorination which comprises fluorinating a monosaccharide or monosaccharide bonded to a base of a nucleic acid by bringing the monosaccharide or monosaccharide bonded to a base of a nucleic acid and a fluorinating agent represented by the general formula (I):




wherein Y represents nitrogen atom or phosphorus atom, R⁰, R¹ and R² represent hydrogen atom or an alkyl or aryl group which may have substituents, the atom and the groups represented by R⁰, R¹ and R² may be a same with or different from each other, and two or three of the groups represented by R⁰, R¹ and R² may be bonded to each other to form a ring into reaction with each other under irradiation with at least one of microwave and electromagnetic wave having a wavelength around a microwave region.

5. (Currently amended) A method of fluorination according to Claim 4, wherein the monosaccharide or monosaccharide bonded to a base of a nucleic acid is fluorinated by bringing the monosaccharide or monosaccharide bonded to a base of a nucleic acid and the fluorinating agent represented by general formula (I):




wherein Y represents nitrogen atom or phosphorus atom, R^0 , R^1 and R^2 represent hydrogen atom or an alkyl or aryl group which may have substituents, the atom and the groups represented by R^0 , R^1 and R^2 may be a same with or different from each other, and two or three of the groups represented by R^0 , R^1 and R^2 may be bonded to each other to form a ring into reaction with each other under irradiation with microwave having a frequency of 1 to 30 GHz.

6. (Previously presented) A method of fluorination according to Claim 4, wherein the fluorinating agent is a compound represented by general formula (II):



wherein Y represents nitrogen atom or phosphorus atom, X represents hydrogen atom or a halogen atom, R^0 , R^1 and R^2 represent hydrogen atom or an alkyl or aryl group which may have substituents, the atom and the groups represented by R^0 , R^1 and R^2 may be a same with or different from each other, and two or three of the groups represented by R^0 , R^1 and R^2 may be bonded to each other to form a ring.

7. (Original) A method of fluorination according to Claim 6, wherein the fluorinating agent is a compound represented by general formula (III):



wherein R^3 , R^4 and R^5 each independently represent an alkyl or aryl group which may have substituents, X represents hydrogen atom or a halogen atom, and two or three of the groups represented by R^3 , R^4 and R^5 may be bonded to each other to form a cyclic structure.

8. (Original) A method of fluorination according to Claim 7, wherein, in general formula (III), R^3 represents an aryl group which may have substituents, X represents fluorine atom, and R^4 and R^5 represent an alkyl or aryl group having 1 to 32 carbon atoms which may have substituents.

9.-12. (Cancelled).

13. (Previously presented) A method of fluorination according to Claim 6, wherein the fluorinating agent is a compound represented by general formula (II) in which X represents fluorine atom.

14. (Previously presented) A method of fluorination according to Claim 6, wherein the fluorinating agent is a compound represented by general formula (II) in


which X represents fluorine atom, Y represents nitrogen atom, R^0 represents 3-methylphenyl group or 2-methoxyphenyl group, and R^1 and R^2 represent ethyl group.

15.-20. (Cancelled).

21. (Previously presented) A method of fluorination according to Claim 4, wherein the fluorination is conducted in a presence of an agent accelerating a reaction.

22.-23. (Cancelled).

24. (Previously presented) A method of fluorination according to Claim 5, wherein the fluorinating agent is a compound represented by general formula (II):



(II)

wherein Y represents nitrogen atom or phosphorus atom, X represents hydrogen atom or a halogen atom, R^0 , R^1 and R^2 represent hydrogen atom or an alkyl or aryl group which may have substituents, the atom and the groups represented by R^0 , R^1 and R^2 may be a same with or different from each other, and two or three of the groups represented by R^0 , R^1 and R^2 may be bonded to each other to form a ring.

25. (Previously presented) The method of fluorination according to Claim 1, wherein the monosaccharide is selected from the group consisting of glucose,

fucose, N-acetylglucosamine, N-acetylgalactosamine, N-acetylneuraminic acid, erythrose, threose, ribose, arabinose, xylose, alose, lyxose, altrose, mannose, gulose, idose, galactose, talose, psicose, fructose, sorbose, tagatose, hexanose, apiose, and a deoxy sugar, an amino sugar, a thio sugar, a condensed sugar and an anhydride of the monosaccharide, and the monosaccharide bonded to a base of a nucleic acid is selected from the group consisting of a nucleoside, an oligonucleoside, ribonucleic acid and deoxyribonucleic acid.